

# Macro II: Stochastic Processes I

Professor Griffy

UAlbany

Spring 2022

# Introduction

- ▶ Today: start talking about time series/stochastic processes.
- ▶ Homework due on Thursday.
- ▶ Continue stochastic processes on Thursday.

# Stochastic Processes

- ▶ Random variables
- ▶ Conditional distributions
- ▶ Markov processes

## Preliminaries

- ▶  $X$  is a random variable,  $x$  is its realization
- ▶ Support: smallest set  $S$  such that  $\Pr(x \in S) = 1$
- ▶ Cumulative distribution function:  $F(x) = \Pr(X \leq x)$
- ▶ Density function:  $f(x) = \frac{d}{dx}F(x)$  implying that  $f(x) dx = dF(x)$

# The Expected Value

- ▶ Mean is the expectation

$$\bar{X} = E(X) = \int_{-\infty}^{\infty} x dF(x) = \int_{-\infty}^{\infty} x f(x) dx$$

- ▶ The expectation of a function of a random variable,  $g(X)$ , is

$$E(g(X)) = \int_{-\infty}^{\infty} g(X) dF(x)$$

- ▶ Note that  $E(g(X)) \neq g(\bar{X})$  unless  $g(X)$  is linear, i.e.

$$g(X) = b \cdot X$$

# The Variance

- ▶ Variance

$$V(X) = E[(X - \bar{X})^2]$$

- ▶ Standard deviation

$$[V(X)]^{\frac{1}{2}}$$

# Jointly Distributed Random Variables

- ▶ Random vector  $(X, Y)$
- ▶ Joint distribution function:  $F(x, y) = \Pr(X \leq x, Y \leq y)$
- ▶ Covariance:  $C(X, Y) = E[(X - \bar{X}) \cdot (Y - \bar{Y})]$
- ▶ Cross-correlation  $= \frac{C(X, Y)}{[V(X) \cdot V(Y)]^{\frac{1}{2}}}$
- ▶ Expectation of a linear combination

$$E(aX + bY) = aE(X) + bE(Y)$$

# What is a Stochastic Process?

- ▶ Stochastic process is an infinite sequence of random variables  $\{X_t\}_{t=-\infty}^{\infty}$
- ▶ j'th autocovariance =  $\gamma_j = C(X_t, X_{t-j})$
- ▶ Strict stationarity: distribution of  $(X_t, X_{t+j_1}, X_{t+j_2}, \dots, X_{t+j_n}, )$  does not depend on  $t$
- ▶ Covariance stationarity:  $\bar{X}_t$  and  $C(X_t, X_{t-j})$  do not depend on  $t$



## Defining a Conditional Density

- ▶ Work with random vector  $\underline{x} = (X, Y) \sim F(x, y)$ .
  - ▶  $X$  and  $Y$  are random variables
  - ▶  $x$  and  $y$  are realizations of the random variables
  - ▶  $F(x, y)$  is joint cumulative distribution
  - ▶  $f(x, y)$  is joint density function

# Conditional Variables and Independence

- ▶ Conditional probability
  - ▶ when  $\Pr(\underline{x} \in B) > 0$ ,

$$\Pr(\underline{x} \in A | \underline{x} \in B) = \Pr(A|B) = \frac{\Pr(A \cap B)}{\Pr(B)}.$$

- ▶ Conditional distribution  $F(y|x)$  (handles  $\Pr(B) = 0$ )
  - ▶ Marginal distribution:  $F_X(x) = \Pr(X \leq x)$
  - ▶  $F(y|x)$  is  $\Pr(Y \leq y)$  conditional on  $X \leq x$

## Defining a Conditional Density

- ▶ Independence: The random variables  $X$  and  $Y$  are independent if

$$F(x, y) = F_X(x) F_Y(y)$$

- ▶ If  $X$  and  $Y$  are independent, then

$$F(y|x) = F_Y(y)$$

and

$$F(x|y) = F_X(x)$$

- ▶ i.i.d means independent and identically distributed
- ▶ Conditional (mathematical, rational) expectation

$$E(Y|x) = \int_{-\infty}^{\infty} y dF(y|x) = \int_{-\infty}^{\infty} y f(y|x) dy.$$

# Markov Property

- ▶ A particular conditional process is called a Markov chain.
- ▶ Markov Property: A stochastic process  $\{x_t\}$  is said to have the Markov property if for all  $k \geq 1$  and all  $t$ ,

$$Prob(x_{t+1}|x_t, x_{t-1}, \dots, x_{t-k}) = Prob(x_{t+1}|x_t) \quad (1)$$

- ▶ That is, the dependence between random events can be summarized exclusively with the previous event.
- ▶ This allows us to characterize this process with a Markov chain.
- ▶ Markov chains are a key way of characterizing stochastic events in our models.

# Markov Chains

- ▶ For a stochastic process with the Markov property, we can characterize the process with a Markov chain.
- ▶ A time-invariant Markov chain is defined by the tuple:
  1. an  $n$ -dimensional state space of vectors  $e_i, i = 1, \dots, n$ ,
    - ▶ where  $e_i$  is an  $n \times 1$  vector where
    - ▶ the  $i$ th entry equals 1 and the vector contains 0s otherwise.
  2. a transition matrix  $P$  ( $n \times n$ ), which records the conditional probability of transitioning between states
  3. a vector  $\pi_0$  ( $n \times 1$ ), that records the unconditional probability of being in state  $i$  at time 0.
- ▶ The key object here is  $P$ . Elements of this matrix are given by

$$P_{ij} = \text{Prob}(x_{t+1} = e_j | x_t = e_i) \quad (2)$$

- ▶ In other words, if you're in state  $i$ , this is the probability you enter state  $j$ .

# Markov Chains

- ▶ Some assumptions on  $P$  and  $\pi_0$ :

- ▶ For  $i = 1, \dots, n$ ,  $P$  satisfies

$$\sum_{j=1}^n P_{ij} = 1 \quad (3)$$

- ▶  $\pi_0$  satisfies

$$\sum_{i=1}^n \pi_{0i} = 1 \quad (4)$$

- ▶ Where does this first property become useful?
- ▶ How would you calculate  $\text{Prob}(x_{t+2} = e_j | x_t = e_i)$ ?

$$= \sum_{h=1}^n \text{Prob}(x_{t+2} = e_j | x_{t+1} = e_h) \text{Prob}(x_{t+1} = e_h | x_t = e_i) \quad (5)$$

$$= \sum_{h=1}^n P_{ih} P_{hj} = P_{ij}^{(2)} \quad (6)$$

# Markov Chains

- ▶ This is also true in general:

$$\text{Prob}(x_{t+k} = e_j | x_t = e_i) = P_{ij}^{(k)} \quad (7)$$

- ▶ Why is this useful? We can use  $\pi_0$  with this transition matrix to characterize the probability distribution over time:

$$\pi_1' = \pi_0' P \quad (8)$$

$$\pi_2' = \pi_0' P^2 \quad (9)$$

$$\vdots \quad (10)$$

- ▶ Thus, by knowing the initial distribution and the transition matrix,  $P$ , we know the distribution at time  $t$

## Stationary Distributions

- ▶ Where does this trend to over time?
- ▶ We know that the transition of the distribution takes the form  $\pi'_{t+1} = \pi'_t P$ .
- ▶ This distribution is stationary if

$$\pi_{t+1} = \pi_t \tag{11}$$

- ▶ (we will relax this to  $t$  large enough momentarily)
- ▶ This means that for a stationary distribution,  $\pi, P$  satisfy

$$\pi' = \pi' P \text{ or} \tag{12}$$

$$(I - P')\pi = 0 \tag{13}$$

- ▶ Anyone recognize this?



## Stationary Distributions

$$\pi' = \pi' P \text{ or} \quad (14)$$

$$(I - P')\pi = 0 \quad (15)$$

- ▶ It is useful to note (and will be useful when we think of linearized solution techniques), that
  - ▶  $\pi$  is the (normalized) eigenvector of the stochastic matrix  $P$ .
  - ▶ In this case, the eigenvalue (root) is 1.
- ▶ A lot of linearizing dynamic systems is about
  - ▶ finding eigenvectors with corresponding eigenvalues of less than 1 (non-explosive).
  - ▶ solving for initial conditions that are orthogonal to the explosive eigenvectors (i.e., the system does not explode).

## Asymptotically Stationary Distributions

- ▶ What about when  $\pi_0 \neq \pi_t$ ? Can it still have a notion of stationarity?
- ▶ Yes. Asymptotic stationarity.
- ▶ Asymptotic stationarity:

$$\lim_{t \rightarrow \infty} \pi_t = \pi_\infty \quad (16)$$

- ▶ where  $\pi'_\infty = \pi'_\infty P$
- ▶ Next, is this ending point unique?
- ▶ Does  $\pi_\infty$  depend on  $\pi_0$ ?
- ▶ If not,  $\pi_\infty$  is an invariant or stationary distribution of  $P$ .
- ▶ This will be very useful when we talk about heterogeneous agents.

## Some Examples

- ▶ Let's pick a simple initial condition:  $\pi'_0 = [1 \ 0 \ 0]$ .
- ▶ And a matrix

$$P = \begin{bmatrix} 0.9 & 0.1 & 0 \\ 0.2 & 0.6 & 0.2 \\ 0.1 & 0.2 & 0.7 \end{bmatrix} \quad (17)$$

- ▶ Now use Matlab to iterate.

# Preliminaries

```
>> piMat = MMat'*piMat  
  
piMat =  
  
    0.9000  
    0.1000  
         0
```

Figure: First iteration

```
>> piMat = MMat'*piMat  
  
piMat =  
  
    0.8300  
    0.1500  
    0.0200
```

Figure: 2nd iteration

```
>> piMat = MMat^(100)*piMat  
  
piMat =  
  
    0.6154  
    0.2308  
    0.1538
```

Figure: First iteration

```
>> piMat = MMat^(1000)*piMat  
  
piMat =  
  
    0.6154  
    0.2308  
    0.1538
```

Figure: Grid of  $k$  values

- ▶ This distribution ( $P$ ) is asymptotically stationary!
- ▶ Unique? Try  $\pi'_0 = [0 \ 0 \ 1]$

# Preliminaries

```
>> piMat = MMat'*piMat  
  
piMat =  
  
    0.9000  
    0.1000  
         0
```

Figure: First iteration

```
>> piMat = MMat'*piMat  
  
piMat =  
  
    0.8300  
    0.1500  
    0.0200
```

Figure: 2nd iteration

```
>> piMat = MMat^(100)*piMat  
  
piMat =  
  
    0.6154  
    0.2308  
    0.1538
```

Figure: First iteration

```
>> piMat = MMat^(1000)*piMat  
  
piMat =  
  
    0.6154  
    0.2308  
    0.1538
```

Figure: Grid of  $k$  values

- ▶ This distribution ( $P$ ) is (probably) a unique invariant distribution.
- ▶ How would we prove this?

# Ergodicity

- ▶ We would like to be able to replace conditional expectations with unconditional expectations.
- ▶ i.e., not indexed by time or initial conditions.
- ▶ Some preliminaries:
  - ▶ Invariant function: “a random variable  $y_t = \bar{y}'x_t$  is said to be invariant if  $y_t = y_0$ ,  $t \geq 0$ , for all realizations of  $x_t$ ,  $t \geq 0$  that occur with positive probability under  $(P, \pi)$ .”
- ▶ i.e., the state  $x$  can move around, but the outcome  $y_t$  stays constant at  $y_0$ .

# Ergodicity

- ▶ Ergodicity:
  - ▶ “Let  $(P, \pi)$  be a stationary Markov chain. The chain is said to be ergodic if the only invariant functions  $\bar{y}$  are constant with probability 1 under the stationary unconditional probability distribution  $\pi$ .”
- ▶ In other words, for any initial distribution, the only functions that satisfy the definition of an invariant function are the same.

## Next Time

- ▶ More stochastic processes.
- ▶ Homework due Thursday.